

Examining Processed Foods in Food Banks and the Presence of Chronic Diseases in
Food Bank Clientele

by

Elizabeth Counts

A thesis presented to the Honors College of Middle Tennessee State University in
partial fulfillment of the requirements for graduation from the University Honors
College

Fall 2024

Thesis Committee:

Dr. Elizabeth A Smith, Thesis Director

Dr. Joan E. McRae, Thesis Committee Chair

Examining Processed Foods in Food Banks and the Presence of Chronic Diseases in Food
Bank Clientele

by Elizabeth Counts

APPROVED:

Dr. Elizabeth A Smith RD, LDN, Thesis Director,
Associate Professor, Department of Human Sciences

Dr. Joan E. McRae, Thesis Committee Chair
Professor, Department of World Languages, Literatures,
and Cultures

Dedication

I would like to dedicate this thesis to my mom and dad whose hearts are full of generosity, kindness, and compassion for others. I will forever admire them and be grateful for what they have instilled in me.

Acknowledgements

To my thesis director, Dr. Smith, thank you for your continuous encouragement, support and guidance. This project would not have been possible without you. I would like to thank Nourish Food Bank and all their staff for so graciously allowing me to conduct my research at their facility. I would also like to thank the clients of Nourish for their willingness to participate in my survey, the kindness they showed, and for sharing their stories with me. To the MTSU Honors College, thank you for giving me the opportunity to explore this topic and complete a project that I am so passionate about. Lastly, I would like to express my deepest gratitude to my friends and family for their belief in me and for their continued support and encourage in everything that I do.

Abstract

Food insecurity affects millions of Americans and puts them at an increased risk of developing chronic disease. Food banks are essential in improving food access for this population. However, previous studies have identified gaps in the nutritional quality of items donated to food banks, thus decreasing clients' access to nutritionally adequate food. The first part of this study aimed to examine the presence of chronic disease in food bank clientele by distributing a survey to clients in the waiting room of a local food bank. The second part assessed the nutritional quality of donated items based on their level of processing using the NOVA Food Classification tool. The study found that the majority of food bank clients had been diagnosed with a diet-related chronic disease, and that slightly over half of donations were ultra-processed foods. This information can be used to form collaborative approaches between food bank staff, their clients, community leaders, and donors to help improve access to nutritious food for those facing food insecurity.

Table of Contents

Dedication.....	iii
Acknowledgements.....	iv
Abstract.....	v
Introduction.....	1
Methodology I.....	4-5
Results I.....	5-6
Figure 1: Food Pantry Clients versus U.S. Population.....	6
Methodology II.....	6-8
Results II.....	8
Figure 2: Food Bank Donations.....	9
Discussion.....	9-13
Limitations and Future Research.....	10
Implications.....	11
Conclusion.....	13-14
References.....	15-18
Appendix A.....	19-24
IRB Approval Letter.....	25

Introduction

Food security is defined as having access to enough food for an active, healthy life.

Conversely, food insecurity is defined as a household-level economic and social condition of limited or uncertain access to adequate food.¹ Sixty-three thousand food pantries across the U.S. provide food for millions of Americans facing food insecurity.¹ However, due to the prevalence of chronic diseases among this population, it is important to consider the role that nutrition plays in this issue.

According to the United States Department of Agriculture, food insecurity affects 44.2 million Americans.¹ Minority populations are disproportionately affected. Black and Hispanic populations are the most at risk of becoming food insecure compared to non-Hispanic whites. Households with children are also more likely to face food insecurity, especially single parent households. Additionally, food insecurity among older adults living alone and households with no children has been increasing.¹ There are several factors that may lead to a person or family becoming food insecure, such as poverty, loss of a job, lack of affordable housing, chronic health conditions making it difficult to work, medical emergencies, and natural disasters.²

Food insecurity has major health impacts on those it affects. A study published by the USDA analyzing five years' worth of data from the National Health Interview Survey found that food insecurity is associated with ten chronic diseases including hypertension, coronary heart disease, cancer, diabetes, and kidney disease.³ Researchers found that that the number of chronic diseases in low food-security households was 18% higher than in high food-security households. They found that high blood pressure, for example, was 10.5 percentage points more likely in adults with very low food-insecurity than in those

with high food-security.³ Another study on food insecurity and chronic disease had similar findings, concluding that food insecurity was strongly associated with hypertension and diabetes. They found that that the risk of diabetes was 50% higher in food-insecure adults than in adults in food-secure households.⁴ Not only do these conditions result in negative physical and mental health outcomes, but they can also lead to further economic hardship with increased doctor visits, hospital stays, and medications payments.

Many who are facing food insecurity receive food from food assistance programs such as food pantries, soup kitchens, and shelters. An estimated 1 in 6 Americans rely on these facilities.² Food pantries provide food for 90% of those who rely on food assistance programs while soup kitchens and shelters make up the remaining amount. Food pantries distribute food directly to individuals from larger warehouse-like facilities called food banks. Food banks receive food from three sources.⁵ The first is from the United States Department of Agriculture (USDA) which purchases food from farmers and then distributes it to food banks. The second source is food purchased by food banks themselves with monetary donations. However, the largest source of food comes from donations from individuals, companies, and local businesses.⁵ Due to the constraints surrounding collecting, storing, and distributing donated food, such as lack of refrigeration, the majority of these foods tend to be canned or packaged goods that can maintain a long shelf life. These foods are often energy-dense and solve the more pressing issue of hunger. However, the nutritional content and the effects of long-term consumption of these types of packaged foods is important to consider, especially for populations at risk of developing or already diagnosed with chronic disease.

Food pantries are essential organizations for fighting food insecurity and it is of primary importance to address this issue of hunger. However, it is also important that these populations are not only given access to food, but to food that provides health-promoting nutrients and that are limited in additives that may put them at risk of developing diet-related chronic diseases. Several studies have investigated food banks to identify the predominant food or nutrient groups in their supplies and those that are in short supply. One study examining several food pantries in Massachusetts found that the largest supplied groups of food were fats, oils, and sweets followed by breads. They also found that vitamin C and calcium were deficient in the food provided by the pantries.⁶ Another study found that, after excluding sweets and fats, the bread, cereal, rice, and pasta groups were the largest while the pantries tended to be limited in food groups such as dairy and vegetables. They also found that over a one-month period 19,590 sweets and 12,041 servings of fat were distributed each day which they noted primarily came from donations of candy, desserts, and cream cheese.⁵

Few studies, however, have looked specifically at processed foods in food pantries. According to Harvard Health, “unprocessed or minimally processed foods are whole foods in which the vitamins and nutrients are still intact. The food is in its natural (or nearly natural) state” while processed and ultra-processed foods are foods that have added ingredients such as sugar, salt, fat, and artificial colors or preservatives.⁷ These products are often high in refined sugars, sodium, saturated fats, and trans fats while being low in fiber and other nutrients. Processed food has been linked to various chronic diseases. A systematic review of 38 studies concluded that there is overwhelming evidence that a higher consumption of ultra-processed foods is positively associated with

obesity and diet-related diseases such as high blood pressure and diabetes mellitus.⁸ This study aims to examine chronic disease prevalence in food pantry clientele and explore the proportion of ultra-processed food present in food pantries. It was hypothesized that there would be a high percentage of food pantry clientele with one or more chronic diseases as well as a high proportion of ultra-processed food available to these clients.

Methodology Part I

The purpose of Part I of the research was to determine how many clients of Nourish Food Bank have one or more diet-related chronic disease. To accomplish this, a Qualtrics© survey was administered to clients while they waited in the Nourish waiting room. Data was collected over a one-month period to prevent any duplication, as clients are permitted to visit Nourish Food Bank for provisions once every thirty days. Nourish operates Monday through Friday and survey results were collected throughout every day of the week to ensure a widely representative sample. The survey began with a consent page that clients had to agree to before advancing on to the survey questions. The survey contained five questions and took participants no more than two minutes to complete. The client was asked to indicate their age group and gender. Then they were asked to check a box next to any of the listed chronic diseases in which they had been diagnosed. Another question followed asking the client to indicate if a member in their household had been diagnosed with any of the listed chronic diseases. All diseases were listed in Layman terms and included high blood pressure, high blood sugar, high cholesterol, overweight, obese, kidney disease, and any type of cancer. The data collected on chronic disease was then compared to the overall U.S. population using data from the National Center of Health Statistics. Lastly, participants were asked to indicate how often they use

a food pantry. Participants excluded from the survey were those below the age of eighteen and pregnant women.

Results Part I

A total of 137 survey responses were collected. Twenty-eight percent of participants were in the 55 to 64 age range followed by the 25 to 34 age range at 20%. The smallest age group was 18 to 24 at only 3%. Females made up the majority of food pantry clientele at 66% compared to men at 31%. The majority of participants reported having one or more diet-related conditions at 70%. Of those clients, the most reported conditions were hypertension (71%), being overweight or obese (48%), high cholesterol (35%), and diabetes (32%). Others reported having high blood sugar (21%), heart disease (20%), kidney disease (9%), cancer diagnosis (9%), and history of a stroke (7%). Participants also reported that members of their household had diet-related conditions at 40%. Of those household members, the majority had hypertension at 69%. The next most prevalent conditions were overweight or obesity (53%), high cholesterol (44%), diabetes (35%), and high blood sugar (31%). Following that were cancer diagnoses (20%), heart disease (16%), history of stroke (16%), and kidney disease a (13%). The majority of participants (53%) indicated that they visit a food pantry once every month. Other participants visited less frequently with 22% using a food pantry a few times a year and 17% using one every few months.

As displayed in Figure 1, food bank clients were more likely to have high blood pressure at 49.64% compared to 24.8% in the U.S population, representing a 24.8 percentage point difference.⁹ Food bank clients were also more likely to have diabetes (by 13.13 percentage points), heart disease (by 2.67 percentage points), stroke (by 4.37

percentage points), and cancer (by 1.08 percentage points).^{9,10,11} Conversely, food bank clients were less likely to be overweight or obese (by 31.52 percentage points).¹² High cholesterol and high blood glucose were unable to be compared due to the lack of data for the U.S. population.




Figure 1. Chronic disease in food pantry clients compared to the general U.S. population

Methodology II

The purpose of the second part of the research was to determine the health quality of food donations that food pantry clients can access. To accomplish this the NOVA Food Classification System was employed.¹³ This system separates food into four groups based on their level of processing. Group 1 is comprised of unprocessed or minimally processed foods. Unprocessed means that the food is “obtained directly from plants or animals and does not undergo any alteration following their removal from nature.”¹³

Minimally processed is defined as a natural food that has had inedible parts removed or has been dried, frozen, pasteurized, or fermented but has no oils, sugar, salt, or other substances added to it. Examples of Group 1 foods include frozen vegetables, packaged grains such as rice, oats, and corn kernels, eggs, dried fruits, beans, nuts and seeds, milk, and yogurt. Group 2 comprises processed culinary ingredients, which are products extracted from natural foods for cooking or seasoning. These foods include oils made from seeds and nuts, cane or beet sugar, honey, maple syrup, butter, coconut fat, and salt. Group 3 are processed foods which are “products manufactured by industry with the use of salt, sugar, oil or other substances (Group 2) added to natural or minimally processed foods (Group 1) to preserve or to make them more palatable.”¹³ This includes canned vegetables and legumes preserved with salt, fruits in sugar-added syrups, beef jerky, salted or sugared nuts, canned fish, salted, dried, smoked or cured meat, cheeses, and bread. Lastly, Group 4 is ultra-processed foods which are “industrial formulations made entirely or mostly from substances extracted from foods (oils, fats, sugar, starch, and proteins), derived from food constituents (hydrogenated fats and modified starch), or synthesized in laboratories from food substrates or other organic sources (flavor enhancers, colors, and several food additives used to make the product hyper-palatable).”¹³ Examples of these foods include packaged snacks, soda, candy, instant soups/noodles, pre-prepared pizza, chicken nuggets, sausages, hamburger buns, breakfast cereals, pastries, and cakes.¹³

To determine how many foods make up each of the four NOVA categories, an inventory was obtained of all the food donated to Nourish Food Bank. The Nova Food Classification guidelines were used to sort products into unprocessed or minimally

processed (Group 1), culinary ingredients (Group 2), processed (Group 3), and ultra-processed (Group 4) to determine the number of products that make up each group. In cases where the product listed was too vague to determine the exact NOVA group, the ingredients of Walmart Inc.'s Great Value branded versions were assessed. Being the largest supermarket chain in the United States, this provided the most likely ingredients contained in the donated product.¹⁴ Baby formula and baby food was excluded from the inventory and accounted for 160 items. Indecipherable products such as "Kroger box", "Food Lion Box" and "treat bags" were also excluded and likely contained an assortment of food products from any of the four NOVA categories.

Results II

From the inventory of donated products over a one-year period, 44,724 products were assessed.¹⁵ Of those products 51.25% made up the ultra-processed Nova Group 4.¹⁵ Examples of contributors to this group included juices made from concentrates and other sweetened beverages, pop tarts, packaged ramen, canned soups, cereal and protein bars, packaged mac and cheese, instant oatmeal, cracker, and chips.¹⁵ The second largest group was the processed food group (Nova Group 3) with 37.93% of products belonging to it. Some of these products included apple sauce, salted nuts, canned vegetables and fruits with added salt or sugar, canned fish, popcorn, pasta sauce, and canned beans.¹⁵ The third largest food group was unprocessed or minimally processed foods (Nova Group 1) at 10.32%. Examples of these foods in the inventory included fruit, milk, vegetables, frozen meats, beans, and grains.¹⁵ The smallest food group was culinary ingredients (Nova Group 2) at 0.499%. Primary contributors of this group included cooking oils, salt, sugar, and baking powders.¹⁵




Figure 2. Nourish food donations compared by NOVA categories

Discussion

Food insecurity is faced by millions of Americans and has been associated with the development of various diet-related chronic diseases. Many rely on food-assistance programs such as food pantries and food banks. Previous studies have determined gaps in the nutritional quality of donated food bank items including a lack of Vitamin C and calcium as well as limited donations of vegetable and dairy items⁶. Few studies, however, have examined food donations using the NOVA Food Classification System. The aim of this study was to utilize the NOVA ranking tool in order to determine what proportion of donated products are ultra-processed foods which have been heavily linked to the development of chronic disease. The study also aimed to determine if the same clients consuming these products were likely to be diagnosed with a chronic disease. The findings of this study provide more insight into the characteristics of this population as well as the food that they may have access to.

Of the food bank clients surveyed, it was determined that the majority had been diagnosed with one or more chronic diseases. Over a third of clients had members of their household who had been diagnosed with a chronic disease as well. Compared to the U.S. population, the data showed that these food bank clients were more likely to have high blood pressure, diabetes, heart disease, stroke, and cancer. This builds on previous studies that have linked food insecurity with an increased risk of developing disease.³ This data demonstrates a need for improving factors that contribute to disease risk and worsen conditions that already exist. A major factor is adequate nutrition and limiting the consumption of ultra-processed foods.⁸

The food that these clients have access to was categorized into the four NOVA classification groups including minimally processed, culinary ingredients, processed, and ultra-processed. It was determined that slightly over half of all donated products were ultra-processed. This adds to the previously existing data on nutrition quality gaps in food pantry donations.⁶ Ultra-processed foods tend to be high in calories, sugar, sodium, saturated and trans fats which contributes to their long shelf-life and hyper palatability, but also to their link to an increased risk of chronic disease.⁸ Ultra-processed foods are also lower in fiber, vitamins, minerals, and other nutrients that help protect against disease.⁸ The results showed that the majority of donations are ultra-processed which limits the nutritional quality of food that clients have access to. The fact that the majority of clients were diagnosed with a chronic disease emphasizes the need to improve their access to more nutritious food options.

Study Limitations

It should be noted that the data collected for the frequency in which clients use food banks may not be representative of most food bank clientele. The specific food pantry where the study was conducted only allowed clients food provisions every 30 days. Future studies should aim to investigate a wider range of facilities. Another limitation is that the inventory used did not include exact brand names or versions, meaning the ingredients within the product can vary slightly and therefore possibly change their assigned NOVA category. To address this issue, future research could gather more comprehensive descriptions of food products by monitoring donations over an extended period.

Lastly, while the results from this study provide valuable insights to both food pantry clientele and food bank nutrition quality, they should be interpreted with caution due to the lack of data analysis. Both the data on chronic disease collected from this study's survey as well as the survey from the NCHS contained multiple-response questions, inhibiting the ability to conduct statistical testing due to the lack of mutually exclusive data points. Future studies should aim to determine if the differences in chronic disease between food pantry clients and the U.S. population are statistically significant. Additionally, in order to add to the data collected on the proportion of donations in each of the four NOVA categories, more extensive future studies should investigate how these proportions compare to those of an average American diet, or the products supplied in an average U.S grocery store.

Implications

As stated previously, the largest source of food supplied at food banks comes from generous donations made by individuals and companies. There are several

challenges in increasing the health quality of these donated items. The first is simply that most donators may not have knowledge of the issue that this study aims to bring awareness to. Another challenge is that donations are likely influenced by the food consumption habits and products available to the U.S. population, a large characteristic of which is processed and ultra-processed foods.¹⁶ This represents a much larger issue in our food system that cannot be easily fixed. Another challenge is the limited refrigeration and freezer storage of food bank facilities and, in some cases, their clients. This makes processed and ultra-processed foods ideal due to their ability to maintain a long shelf-life. Nevertheless, there are still solutions that could improve the health quality of donated items.

One possible solution could be a resource that provides guidance for those purchasing items to donate. This list could include healthier versions of shelf-stable products such as low sodium canned soups and vegetables, no sugar added fruits, dried grains, canned fish, dried or canned legumes, nuts, popcorn, peanut butter, and low-sugar cereals. Some food banks have moved a step further and implemented nutrition quality regulatory policies, eliminating the distribution of items like candy and soda. One study gathering the opinions of food bank directors and staff showed varying responses to this.¹⁷ Many were concerned that this would harm donator relationships thus decreasing overall donations. However, a few food banks who have implemented these policies on soda donations found that donors were willing to accommodate this request and in cases where some donators were lost, more were gained to fill in this gap. The study noted that more research is needed to determine if regulatory nutritional policies are an effective approach in improving nutritional quality without decreasing donations.¹⁷

Another challenge that food bank staff cited was being able to properly store and distribute perishable goods.¹⁷ This requires adequate refrigeration, staff and volunteers to process these goods, and a means of transporting goods to clients. Encouraging monetary donations to food banks and increasing their resources may help mitigate these barriers and improve the distribution process. For example, many food banks nationwide have been able to implement mobile pantries to quickly distribute perishable items.¹⁷

One solution for obtaining more of these nutrient-dense items is to increase partnerships between food banks and local farmers. These programs already exist in several states such as Connecticut, Georgia, Kentucky, Maryland, Oregon and Rhode Island.¹⁸ Other states like Arizona, California, Colorado, Oregon, Iowa, and Kentucky offer tax credits to farmers donating extra produce to food banks.¹⁸ Furthermore, programs like Grow to Give and Ample Harvest encourage home and community gardeners to donate a portion of their produce to food banks as well.¹⁸

Conclusion

Food insecurity impacts millions of Americans, and those who face food insecurity are at a greater risk of developing chronic disease. This also puts them at a greater risk of further financial instability from medical bills and hospital stays. Many who are food insecure rely on food banks and food pantries which are largely dependent on donations. This means that food banks are limited in their control over the food they supply, and their clients are limited in what food they have access to. Additional research is needed to accurately assess how the prevalence of chronic diseases among food pantry clients compares to that of the U.S. population. However, this study displays valuable insights into the health characteristics of food pantry clientele and the types of items food banks

typically receive. The majority of clients surveyed had been diagnosed with one or more diet-related chronic diseases, and slightly over half of all food bank items were ultra-processed. Information in this study can be used to draw attention to this gap in access to nutritious food combined with the health risks already present in this population. From there, a collaborative approach between food bank staff, their clients, community leaders, and donors can be used to develop solutions to help fill this gap. It is important that those facing food insecurity not only have access to food, but nutritionally adequate food that is essential for maintaining health, preventing disease, and contributing to a higher quality of life.

References

1. Coleman-Jensen A, Reed-Jones M, Hales LJ, Rabbitt MP. Key Statistics & Graphics. USDA ERS - Key Statistics & Graphics. October 25, 2023. Accessed April 1, 2024. <https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-u-s/key-statistics-graphics/>.
2. Feeding America. What is food insecurity? Feeding America. Accessed April 1, 2024. <https://www.feedingamerica.org/hunger-in-america/food-insecurity>.
3. Gregory CA, Coleman-Jensen A. Food insecurity, chronic disease, and health among working-age adults. USDA ERS. July 2017. Accessed April 1, 2024. <https://www.ers.usda.gov/publications/pub-details/?pubid=84466>.
4. Seligman HK, Laraia BA, Kushel MB. Food insecurity is associated with chronic disease among low-income NHANES participants. *The Journal of Nutrition*. 2010;140(2):304-310. doi:10.3945/jn.109.112573
5. Cotugna N, Vickery CE, Glick M. An outcome evaluation of a Food Bank Program. *Journal of the American Dietetic Association*. 1994;94(8):888-890. doi:10.1016/0002-8223(94)92370-1
6. Akobundu UO, Cohen NL, Laus MJ, Schulte MJ, Soussloff MN. Vitamins A and C, calcium, fruit, and dairy products are limited in food pantries. *Journal of the American Dietetic Association*. 2004;104(5):811-813. doi:10.1016/j.jada.2004.03.009
7. Katherine D. McManus M. What are ultra-processed foods and are they bad for our health? Harvard Health. January 9, 2020. Accessed April 1, 2024.

<https://www.health.harvard.edu/blog/what-are-ultra-processed-foods-and-are-they-bad-for-our-health-2020010918605>.

8. Jardim MZ, Costa BV, Pessoa MC, Duarte CK. Ultra-processed foods increase noncommunicable chronic disease risk. *Nutrition Research*. 2021;95:19-34. doi: 10.1016/j.nutres.2021.08.006
9. National Center for Health Statistics. NHIS - tables of summary health statistics: Circulatory Disease. Centers for Disease Control and Prevention. November 5, 2019. Accessed August 10, 2024. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-15.pdf
10. National Center for Health Statistics. NHS - tables of summary health statistics: Diabetes and Kidney Disease. Centers for Disease Control and Prevention. November 5, 2019. Accessed August 10, 2024. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-4.pdf
11. National Center for Health Statistics. NHS - tables of summary health statistics: Cancer. Centers for Disease Control and Prevention. November 5, 2019. Accessed August 10, 2024. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-3.pdf
12. National Center for Health Statistics. NHS - tables of summary health statistics: Overweight and Obese. Centers for Disease Control and Prevention. November 5, 2019. Accessed August 10, 2024.

https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-3.pdf

13. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Jaime P, Martins AP, Canella D, Louzada M, Parra D. NOVA. The star shines bright. Food classification. *World Nutrition*. 1-3, 28-38. <https://archive.wphna.org/wp-content/uploads/2016/01/WN-2016-7-1-3-28-38-Monteiro-Cannon-Levy-et-al-NOVA.pdf>
14. Walmart Inc. FY23 Financial Fact Book. Walmart Annual Reports. April 20, 2023. Accessed September 20, 2024. https://s201.q4cdn.com/262069030/files/doc_downloads/Factbook/2023/FY23-Financial-Fact-Book.pdf.
15. *Food Inventory Database*. Nourish Food Bank Murfreesboro; 2024.
16. Martínez Steele E, Popkin BM, Swinburn B, Monteiro CA. The share of ultra-processed foods and the overall nutritional quality of diets in the US: Evidence from a nationally representative cross-sectional study. 2017;15(1). doi:10.1186/s12963-017-0119-3
17. Handforth B, Hennink M, Schwartz MB. A qualitative study of nutrition-based initiatives at selected food banks in the Feeding America Network. *Journal of the Academy of Nutrition and Dietetics*. 2013;113(3):411-415. doi:10.1016/j.jand.2012.11.001
18. Healthy Food Initiatives in food pantries. County Health Rankings & Roadmaps. December 7, 2020. Accessed August 10, 2024.

https://www.countyhealthrankings.org/strategies-and-solutions/what-works-for-health/strategies/healthy-food-initiatives-in-food-pantries#footnote_25.

Appendix A

Qualtrics© Chronic Disease Survey

Presence of Chronic Disease in Food Pantry Clientele

Q1 You are being asked to participate in a research project. The following information is provided to inform you about the research project and your participation in it.

Study Title: Examining the Correlation Between Processed Foods in Food Pantries and the Presence of Chronic Diseases in Food Pantry Clientele

Approval Date: 4/2024

Study: IRB-FY2024-198

Principal Investigator: Elizabeth Counts

Institution: Middle Tennessee State University

Purpose of the study: The purpose of this study is to investigate a link between processed foods at food pantries and whether the clients of those food pantries have 1 or more chronic diseases.

Procedure: If you consent to participate in this study, you will proceed to a survey where you will check the box next to any chronic diseases you or a family member may have. It will take around 5 minutes to complete.

Compensation for participation: N/A

Here are your rights as a participant:

- a) Your participation in this research is voluntary.
- b) You may skip any item that you don't want to answer, and you may stop the research at any time. Note that if you leave an item blank, you will be warned that you missed one, just in case it was an accident. You can still click that you don't want to answer. Some items may be required in order to accurately present the study.
- c) There are no risks associated with your participation besides possible discomfort with some of the questions. There are no real benefits to you from participating

besides possibly learning something about the research.

d) You will NOT be asked to provide any identifiable personal information.

e) All efforts, within reason, will be made to keep the personal information in your research record private, but total privacy cannot be promised. Your information may be shared with people at MTSU (such as the MTSU Institutional Review Board) or other agencies (such as the Federal Government Office for Human Research Protection) if you or someone else is in danger or if we are required to do so by law.

If you should have any questions about this research study please contact:

Principal Investigator: Elizabeth Counts

Contact Information: 615-806-9342, enc3f@mtmail.mtsu.edu

Faculty Advisor: Dr. Elizabeth Ann Smith

Contact Information: 419-310-1725, Elizabeth.ann.smith@mtsu.edu

For additional information about giving consent or your rights as a participant in this study, please contact the Middle Tennessee State University (MTSU) Office of Compliance at 615-494-8918 or via email at irb_information@mtsu.edu.

If you're ready to get started, please make your choice below before clicking the arrow button. Thanks again for volunteering your time to this project!

I have read the information above. I am at least 18 years old. I believe I understand the purpose, risks, and benefits of the research, and I know what I will be expected to do as a study participant.

I consent

I do not consent

Skip To: End of Survey If Please select your choice below: = I do not consent

Skip To: End of Survey If Please select your choice below: != I consent

Q7 Please select your age group

- 18-24
 - 25-34
 - 35-44
 - 45-54
 - 55-64
 - 65 or older
-

Q6 What is your gender?

- Female
 - Male
 - Other
-

Q2 Select all conditions that apply to you:

- High blood pressure/hypertension
 - High blood sugar
 - Diabetes
 - Heart disease
 - History of a stroke
 - Kidney disease
 - Overweight or obese
 - High cholesterol
 - Diagnosis of any type of cancer (breast, colon, prostate, stomach, pancreatic, ect.)
-

Q3 Select all conditions that apply to a member of your household

- High blood pressure/hypertension
 - High blood sugar
 - Diabetes
 - Heart disease
 - History of a stroke
 - Kidney disease
 - Overweight or obese
 - High cholesterol
 - Diagnosis of any type of cancer (breast, colon, prostate, stomach, pancreatic, ect.)
-

Q4 How often do you visit food pantries?

- Once a week
- Once every 2 weeks
- Once a month
- Once every few months
- A few times each year

End of Block: Default Question Block

IRB Letter

<file:///C:/Users/enc3f/Downloads/IRB%20Approval%20Letter.pdf>