

Chlorine Dioxide Gas: Development, Optimization, and Application, of an Antiviral
Assay Based on the MS2 Bacteriophage

By

Hunter Brady

A thesis presented to the Honors College of Middle Tennessee State
University in partial fulfillment of the requirements for graduation from
the University Honors College

Spring 2023

Thesis Committee:

Dr. Anthony Newsome, Thesis Director

Dr. Stephen Wright, Second Reader

Dr. John DuBois, Thesis Committee Chair

Chlorine Dioxide Gas: Development, Optimization, and Application, of an Antiviral
Assay Based on the MS2 Bacteriophage

by Hunter Brady

APPROVED:

Dr. Anthony Newsome, Thesis Director
Professor, Biology

Dr. Stephen Wright, Second Reader
Professor, Biology

Dr. John DuBois, Thesis Committee Chair
Professor, Biology

Dedication

To my constant life-long supporters, my mother Nikki, father Ron, and brother Zach.

To my brother Austin Duke.

Solus Christus Soli Deo Gloria.

Acknowledgments

First, I express my deepest gratitude to Dr. Anthony Newsome, my research mentor, for his constant support, availability, and guidance. Without Dr. Newsome, this project would not have been possible. I appreciate his steady heart of service, long-suffering, and his love for education.

Additionally, this project would not have been possible without the Office of Research and Sponsored Programs and the Undergraduate Research Center. They have provided not only guidance through partnership but also financial support through the Undergraduate Research Experience and Creative Activity grants that I have received. Specifically, I would like to express my deepest gratitude toward Dr. Jamie Burriss, the program manager of the Undergraduate Research Center, who has constantly supported me personally and through this project.

Lastly, I would like to thank the Middle Tennessee State University Biology Department and my thesis committee: Dr. Anthony Newsome, Thesis Director; Dr. Stephen Wright, Second Reader; and Dr. John DuBois, Thesis Committee Chair.

Abstract

Since emergence of the SARS-CoV-2 virus, the need to identify antiviral agents for disinfection purposes has dramatically increased. Chlorine dioxide gas has previously been identified as an antibacterial agent with strong oxidizing capabilities. The MS2 bacteriophage has previously been identified as a suitable surrogate for the development and application of virucide decontamination methods. The purpose of this study was to identify and assess the antiviral properties of chlorine dioxide gas and to identify optimum physical conditions for potential deployment in support of current antiviral disinfection needs. Using the MS2 bacteriophage model system, preliminary studies used the double-layer agar plaque assay technique to evaluate the antiviral activity of chlorine dioxide gas. Results support the potential use of chlorine dioxide gas as an antiviral agent and that environmental factors heavily influences the ability of chlorine dioxide gas to act as an antiviral agent.

Preface

This undergraduate honors thesis contains results gathered under the direction of Dr. Anthony Newsome at the Middle Tennessee State University Department of Biology. This research was prompted under the need to identify antiviral agents in response to the COVID-19 pandemic in conjunction with Dr. Newsome's previous research interests including chlorine dioxide as an antibacterial agent.

As an undergraduate student, I was interested in research for many years. When I transferred to Middle Tennessee State University, I came across Dr. Anthony Newsome's faculty profile and found his research interests particularly intellectually fascinating. Under his guidance, this research was accomplished. Entering medical school at Lincoln Memorial DeBusk College of Osteopathic Medicine, I hope to continue fostering a life dedicated to pursuing intellectual knowledge and research to build a brighter tomorrow.

Table of Contents

Dedication.....	i
Acknowledgments.....	ii
Abstract.....	iii
Preface.....	iv
Table of Contents.....	v
List of Tables.....	vi
List of Figures.....	vii
CHAPTER I: INTRODUCTION.....	1
Background.....	1
Viruses.....	1
Brief History of Disinfectants.....	3
Research Focus.....	7
CHAPTER II: MATERIALS AND METHODS.....	8
Retrieval and Preparation of Viral and Bacterial Cultures and Media.....	8
Chlorine Dioxide Generation System and Deployment Container.....	9
Preparation of Test Materials.....	9
Treatment Protocol.....	10
Viral Dilution and Plaque Assay.....	10
Mathematical Analysis.....	11
CHAPTER III: RESULTS.....	13
Development of the Testing Protocol.....	13
Antiviral Capabilities of ClO ₂ Gas on Metal.....	13
Antiviral Capabilities of ClO ₂ Gas on Cloth.....	15
CHAPTER IV: DISCUSSION.....	20
Summary of Findings.....	20
Implications.....	21
Bibliography.....	22

List of Tables

Table 1: Investigation of the effects of ClO ₂ gas on MS-2 inoculated stainless-steel coupons at room temperature and ≤ 64% relative humidity	16
Table 2: Investigation of the effects of ClO ₂ gas on MS-2 impregnated cloth at room temperature and ≤ 64%- ≥90 % relative humidity	16

List of Figures

Figure 1: Enveloped Viral anatomy	1
Figure 2: Bacteriophage replication cycle (Cowan et al. 2019).....	2
Figure 3: Common targets of for inactivation by chlorine dioxide	6
Figure 4: Research objectives flow chart.....	7
Figure 5: Example of plaque assay	11
Figure 6: General outline of the methodologies developed and employed in the pursuit of the determination of the antiviral properties of ClO ₂ gas	12
Figure 7: Examination of the effects of wet versus dry viral suspension contaminated stainless steel surfaces exposed to 200 ppm ClO ₂ at room temperature and ≤ 64% humidity for 45 minutes.....	17
Figure 8: Examination of the effects of 1:10 diluted MS2 bacteriophage stock solution with sterile H ₂ O prior to 200 ppm ClO ₂ gas exposure for 45 minutes at room temperature and <64% humidity.....	17
Figure 9: Examination of the antiviral capabilities of ClO ₂ on cloth impregnated with 1:10 diluted MS2 bacteriophage solution at 45 minutes and 960 minutes at room temperature and ≤ 64% relative humidity.....	18
Figure 10: Examination of the antiviral capabilities of ClO ₂ on cloth impregnated with 1:10 diluted MS2 bacteriophage solution at varying concentrations of 200 PPM and 40 ppm ClO ₂ Gas for 45 minutes at room temperature and ≤ 64% relative humidity	18
Figure 11: Examination of the antiviral capabilities of ClO ₂ on cloth impregnated with 90% diluted MS2 bacteriophage solution at varying relative humidity conditions of ≤ 64% and ≥ 90% relative humidity (RH).....	19

CHAPTER I: INTRODUCTION

Background

Since their discovery in the 1800s, viruses have been of interest to the scientific community because of their potential to cause a variety of diseases in humans. Viruses have altered many aspects of people's lives throughout human history (Oldstone 2010). For example, notable diseases caused by viruses include influenza, COVID-19, and human immunodeficiency virus (HIV). It is estimated that nearly 300 million people were killed by smallpox in the twentieth century. In perspective, more than three times more deaths were associated with the virus than wars in the twentieth century (Oldstone 2010). Most recently, the SARS-CoV-2 virus was responsible for the novel outbreak of COVID-19, which has claimed the lives of over 5 million worldwide (Johns Hopkins University 2020). As Nobel Prize winner Peter Medawar says, "it has been well said that a virus is a bad piece of news wrapped up in protein" (Medawar and Medawar 1985). From smallpox to COVID-19, there have been millions of deaths associated with viruses.

Viruses

Viruses are submicroscopic biological agents (Figure 1) and consist of a basic genome (either deoxyribonucleic acid or ribonucleic acid) and a protein coating (Pellett et al. 2014). Categories of viruses are based on structural




Figure 1: Enveloped Viral anatomy
(Bruslind 2018 Feb 6)

features, type of nucleic acid, and the presence or absence of an envelope. Viruses are obligate intercellular parasites, which means that viruses are reliant upon a host cell for replication (Pellett et al. 2014). Intercellular viral replication is a feature of viral infection in humans.

Viral replication can be characterized by absorption, penetration, genomic replication, reassembly, and release (Figure 2) (Cowan et al. 2019). The initial step for viral replication is the protein coating on the virus binding to the host cell's surface; the protein coating and the binding site on the cell are specific (Cowan et al. 2019). The cell must then intake the virus into the cell by endocytosis, which is a packaging of the virus into a vesicle. The virus uses the host cell's environment to replicate the viral genetic information and protein coatings (Cowan et al. 2019). Once the necessary genetic materials are created, the virus reassembles and is released from the cell, completing the replication cycle. Typically, as a result of viral replication, the host cell ruptures and dies.




Figure 2: Bacteriophage replication cycle (Cowan et al. 2019)

Because of the labor required for culture of eukaryotic cell viruses and the pathogenic nature of some viruses, bacteriophages have become an excellent model organism for virological studies (Keen 2015). Bacteriophages are unique viral particles that only infect bacteria (Clokie et al. 2011). Bacteriophages have similar biological features to viruses infecting human eukaryotic cells, thus, allowing bacteriophages to serve as a model organism (Clokie et al. 2011). The bacteriophage MS2, bacteriophage Phi-6, MHV, 229 E, and the SARS-CoV-2 virus were recently compared to evaluate their suitability for virological studies (Ratliff and Oudejans 2021). In comparison, the MS2 bacteriophage offered the most promising potential as a model organism for antiviral studies because of its ease of use and high resistance to inactivation (Ratliff and Oudejans 2021). Additionally, MS2 bacteriophage is easily recognizable because of its well-known conjugate host, *Escherichia coli*.

Brief History of Disinfectants

For thousands of years, the use of disinfectants has been employed to limit the spread of disease. The earliest mention of a chemical disinfectant is noted in 800 B.C. in the Odyssey, mentioning the use of sulfur to disinfect a room (Blancou 1995). In ancient times, sulfur, mercury, copper, acids, and salts were used for disinfection (Blancou 1995). Although it has only been within the last 300 years, disinfectants have gained more widespread use in the scientific community (Hugo 1991). Chlorine, discovered in 1744, became one of the first industrial chemical disinfectants (Hugo 1991). The early 1800s started with advertisements for the use of phenols for disinfection (Virox Technologies).

Later, in 1839, iodine became another common disinfectant used as an antibacterial agent (Hugo 1991).

Additionally, the use of hydrogen peroxide and formaldehyde became widely used by the end of the 19th century (Hugo 1991). The 1950's introduced the use of alcohol-based products for disinfection, and by the 1970's use of ammonium-based products started to be used by commercial manufacturers (Virox Technologies). For almost 3,000 years, humans have employed antimicrobial agents to defend against infectious diseases.

In 1814, Sir Humphrey Davy is reported to be the first person to discover chlorine dioxide (ClO₂), which is now recognized as an antibacterial agent. Davy discovered ClO₂ gas by mixing sulphuric acid with potassium chlorate (Lenntech 2021). Since the discovery, ClO₂ has been used in a variety of ways.

Chlorine dioxide gas and liquid are potent oxidizing agents and have a history of different practical applications. A notable historical use was the use of ClO₂ gas in the disinfection of *Bacillus anthrax* from federal buildings during the intentional release of anthrax spores in 2001 (Rastogi et al. 2010). Chlorine dioxide gas has also historically been used for the disinfection of drinking water (Aieta and Berg 1986). Additionally, ClO₂ has more recently been used in the disinfection of medical equipment such as endoscopes (Isomoto et al. 2006). Although ClO₂ gas and liquid have been historically used for many practical applications, generating the gas or liquid is often challenging.

Previous studies conducted on ClO₂ gas at Middle Tennessee State University have aptly demonstrated the antibacterial properties of the gas. Several studies have examined the potential practical uses of ClO₂ gas (Bhawana et al. 2014; Newsome et al.

2009; Stubblefield and Newsome 2015). Additionally, studies have been aimed at evaluating the use of ClO₂ to inhibit the hemagglutinin protein on the influenza virus (Gormsen 2016). These preliminary studies place Middle Tennessee State University in a unique position to continue investigating the antimicrobial properties of ClO₂.

The use of ClO₂ gas has historically been restrained because of the difficulties in generating the gas. The gas is too unstable for transportation and must be prepared at the application site (Stubblefield and Newsome 2015). Presently, this presents a problem because the gas has typically been generated using trained personnel and expensive machinery (Stubblefield and Newsome 2015). Recently, new technology has emerged, allowing one to quickly produce ClO₂ gas (Smith et al. 2014). This current study is based on use of the ICA-TriNova system, which involves mixing a solid acid and base in a gas-permeable bag to produce ClO₂ gas. It does not require machinery or highly trained professionals to easily produce ClO₂ gas.

Most recently, ClO₂ gas has attracted the attention of scientists because of its oxidative properties that may allow ClO₂ to express antiviral properties. Several studies have been aimed at evaluating the antiviral properties of ClO₂ gas in different model organisms (Ogata et al. 2016; Ogata and Shibata 2008; Ogata 2012; Gormsen 2016; Sanekata et al. 2010). Although most studies published to date evaluate the inactivation of the surface portions of these viruses, they do not address the actual ability of ClO₂ to inactivate the virus. Also, several of the above-cited pieces of literature require that the ClO₂ gas be held at a particular part per million for a specific time for viral inactivation, which is a substantially difficult task (Kály-Kullai et al. 2020). Additionally, much of the literature in circulation have a financial interest in the study's success. Lastly, the current

literature lacks in its ability to show the inactivation of non-enveloped viruses, which are known for their inability to be inactivated in the presence of antiviral agents (Lin et al. 2020). The lack of data to support the ability of ClO₂ gas to inactivate viruses has prompted this current investigation.

Viral targets of ClO₂ are outlined in Figure 3.

A few studies have shown that ClO₂ oxidatively interacts with amino acids, which aid in the infection process by viral particles. Specifically, ClO₂ oxidatively interacts with cysteine, methionine, tyrosine, and tryptophan (Noszticzius et al. 2013).

Studies have also shown that haemagglutinin and neuraminidase are inactivated in the presence of ClO₂ (Ogata and Shibata 2008; Gormsen 2016).

Most studies addressing the mode of viral inactivation by ClO₂ have been investigated using enveloped viruses. In regard to enveloped viruses, one study evaluated the effects of ClO₂ on viral RNA (Alvarez and O'Brien 1982). The study revealed that the positive-sense RNA poliovirus genomic information was affected by ClO₂ (Alvarez and O'Brien 1982). Comprehensive knowledge of the mechanism of action of ClO₂ against viruses remains limited and poorly understood.




Figure 3: Common targets of for inactivation by chlorine dioxide

Research Focus

The objective of this study was to develop a system by which the antiviral activity of ClO₂ gas could be examined, quantified, and analyzed to aid in the investigation of potential antiviral agents (Figure 4). In optimization, factors such as humidity, exposure time to ClO₂ gas, and substrates were examined. Additionally, this study was aimed at developing a system that would be easily recognizable by the Environmental Protection Agency as a valid mechanism for future studies to examine the antiviral properties of ClO₂ gas. This included the use of the MS2 bacteriophage as the model organism of choice due to its high resistance to inactivation as a nonenveloped virus. Thus, studies were modeled in accordance with other Environmental Protection Agency publications and presentations (Oudejans and Ratliff 2021; Wyrzykowska-Ceradini et al. 2019).




Figure 4: Research objectives flow chart

CHAPTER II: MATERIALS AND METHODS

Retrieval and Preparation of Viral and Bacterial Cultures and Media

Viral and bacterial cultures used for this study consisted of the MS2 bacteriophage and *Escherichia coli* as the conjugate host. The MS2 bacteriophage (15597-B1) and *Escherichia coli* (Migula) Castellani and Chalmers (15597) were purchased from American Type Culture Collection (ATCC). Upon receipt of the MS2 bacteriophage, it was reconstituted in 2 ml of deionized water and stored at 4° C. Prior to use, bacteriophage was diluted 1:10 in sterile deionized water and stored at 4° C. Additionally, upon receipt, *Escherichia coli* was cultured and aliquots stored (-70° C). When needed, samples were thawed, plated onto trypticase soy agar (TSA), incubated at 37° C, and stored at room temperature after 24 hours of incubation.

Trypticase soy agar was prepared by mixing 20 grams of TSA powder with 500 mL of deionized purified water, brought to a boil, autoclaved, and evenly divided into petri dishes aseptically. Trypticase soy broth (TSB) was prepared by mixing 15 grams of the powder with 500 mL of deionized purified water, brought to a boil, 4.5 mL was pipetted into test tubes, capped, and autoclaved. Nutrient agar was prepared by mixing 1 gram with 99 mL of deionized purified water, heated to boiling and 4.0 mL were pipetted into test tubes, capped, and autoclaved. All media was stored at 4° C.

Chlorine Dioxide Generation System and Deployment Container

Traditionally, ClO₂ gas has been generated by placing sodium chlorite in aqueous solution causing the production of ClO₂ gas. Although, examination of ClO₂ gas's properties and uses has been difficult due to the gas's instability (Stubblefield and Newsome 2015). Thus, investigation of ClO₂ gas's antiviral properties was modeled on a solid acid base mixture, a novel system of examination. The ICA-Trinova (Newman, GA) ClO₂ gas release system was employed as the deployment mechanism of choice. This novel system, allows scientists to swiftly generate ClO₂ gas in a cost-effective, safe, and easy-to-use way. The system uses two granulated solid chemicals, a base (reagent A) and an acid (reagent B), mixed in a gas permeable pouch to produce ClO₂ gas.

Prior studies conducted at Middle Tennessee State University with ClO₂ have used an air-tight gassing bucket (Gormsen 2016). This bucket has been chosen as a suitable gassing container for the current study due to the airtight lid and internal air circulating fan to promote even distribution of the gas in the chamber. Additionally, for this conducted study, mathematical formulas have been derived by ICA-Trinova for determination of parts per million (PPM) of gas present in the container.

Preparation of Test Materials

A Union 10' x 20" galvanized steel roll was purchased from Lowes and used for the metal treatment protocol. Coupons were cut from the galvanized steel into 15mm x 12 mm coupons, and the edges were bent to create a small well for holding the viral suspension. Metal coupons were then sterilized (via autoclave) and aseptically stored. In

similar fashion, two-ply 100% cotton fabric was obtained and cut into 15 mm x 15 mm slices, sterilized (via autoclave), and aseptically stored.

Treatment Protocol

Treatment using ClO₂ gas was prepared according to ICA-Trinova (Newnan, GA) mixing protocol. Measured equal amounts of reagent A and B were mixed together in a gas permeable bag and hung on a clip inside the gassing container. Substrate of interest (either metal or cloth) was impregnated with MS2 bacteriophage and was placed on top of a test tube rack in the 5 L gassing chamber in which a small battery-operated fan was placed to promote equivalent gas distribution. In subsequent protocols, to ensure humidity stabilization, single use humidity chips (Vetovation Raleigh, NC) were inserted into the testing chamber. Each treatment was performed in duplicate or triplicate along with untreated controls.

Viral Dilution and Plaque Assay

After treatment, the protocol for determining the number of bacteriophages recovered, killed, log-reduction, and percent reduction was based on the traditional viral plaque assay method (Figure 5). This method includes creating dilutions of the stock cultures, mixing the viral particles with the conjugate host cell, and using a double-layer agar technique. This method allows one to quickly quantify the number of viral particles per milliliter by counting the number of plaques present and using mathematical analysis

to establish or determine the amount of virus recovered, killed, and inactivated in treated and untreated controls.

Treated and untreated substrates (containing bacteriophage) were placed in 4.5 mL of TSB and vortexed. The solutions were then diluted ten-fold 5 times in TSB. Recoverable plaque forming units (PFU) were determined by delivery of 0.1 mL of the appropriate dilution into 4.0 mL of 1% nutrient soft agar. To this, 0.1 mL of *E. coli* (in logarithmic growth) was added to the soft agar/phage preparation and immediately poured onto the surface of TSA plates. Plates were incubated (37° C) overnight and recoverable PFU/mL were then determined from the treated and untreated substrate test coupons (Figure 6).



Figure 5: Example of plaque assay

Mathematical Analysis

Mathematical analysis calculations were performed to determine the amount of MS2 bacteriophage recoverable after treatment of a stock culture. For computations, plates with 30-300 plaques were utilized. Plaques were tallied and documented in a

laboratory grade notebook. From the plaque assays, calculations were performed on the plate with the most countable plaques. The number of plaques counted was multiplied by ten (giving the number of recoverable viral particles per mL of that given dilution), then multiplied by ten, the number of times the number on the dilution tube (for example, if the number of plaques on a plate was 250: $250 \times 10 = 2500$ signifying PFU/mL in the 10^{-5} solution, the 2,500 is multiplied by ten, five times giving the number of recoverable viral particles in the stock solution and present from the material exposed to ClO_2). Relative humidity determinations were made using a Thermo-Hygro meter (Fischer Scientific).

1.5-10 microliters of MS-2 bacteriophage was pipetted either onto cloth or metal coupons and was treated with 20-200 ppm of ClO_2 gas

2. Sample was placed into trypticase soy broth, vortexed, and diluted ten-fold

3. Dilutions were plated onto trypticase soy agar and incubated

4. Plaques were counted, and data was recorded




Figure 6: General outline of the methodologies developed and employed in the pursuit of the determination of the antiviral properties of ClO_2 gas

CHAPTER III: RESULTS

Development of the Testing Protocol

In the development of the ClO₂ gas and MS2 bacteriophage testing protocol, preliminary studies were aimed at testing to evaluate if previous experimental protocols would be suitable for the proposed study. This included evaluation of studies published from the Environmental Protection Agency, Middle Tennessee State University, and other laboratories investigating the oxidative properties of ClO₂. All tests were measured against controls that mirrored the protocol given to the test groups except for the exposure to ClO₂ gas. These preliminary testings yielded results that promoted the collection of additional data in support of the antiviral potential of ClO₂.

Antiviral Capabilities of ClO₂ Gas on Metal

Studies were first aimed at the investigation of the antiviral properties of ClO₂ on nonporous surfaces such as stainless-steel metal coupons. The investigation on metal coupons included the testing of 200 PPM ClO₂ gas on small stainless-steel coupons yielding a percent reduction range of 61%-99.99% in recoverable MS2 bacteriophage at 200 PPM for either 45 minutes or 960 minutes (16 hours) gas treatment (Table 1). In this range, it was observed that allowing the MS2 bacteriophage suspension to dry to completion before gas treatment, resulted in a decreased percent reduction with an average of 61% (*SD* = 0.093), while placing the coupon “wet” (not dried suspension) into the gassing chamber yielded a larger percent reduction with an average of 87% (*SD* = 0.024) at 45 minutes (Figure 7).

In optimization of these findings, it was observed that the organic content present in the MS2 bacteriophage suspension can alter the percent reduction of MS2 bacteriophage in the presence of ClO₂ gas. For example, when the stock culture from ATCC was diluted 1:10 in deionized water (thus reducing the organic content by 90%), the percent killed was increased. It was found that dilution of the stock MS2 bacteriophage culture by 10-fold yielded a 99.80% reduction of MS2 bacteriophage after 200 PPM exposure for 45 minutes (Figure 8) and a 99.99% reduction of MS2 bacteriophage after 960 minutes of exposure.

In review of ClO₂ gas antiviral potential using stainless steel coupons, it was found that ClO₂ was capable optimally of acting as an antiviral agent on stainless-steel coupons under certain conditions, such as reduction of organic content in the MS2 bacteriophage suspension and dependent upon the state of the suspension (either evaporated or wet) on the substrate. Log reduction displayed an average of 0.91 log reduction (< 90% reduction) in MS2 bacteriophage on stainless-steel surfaces when the contaminate was immediately exposed to ClO₂ gas, and a 0.41 log reduction in MS2 bacteriophage when the viral suspension was allowed to evaporate to dryness. Additionally, it was found that an average of 2.7 log reduction occurred when the viral suspension was diluted by 90% in water (thus reducing the organic content of the suspension) (Figure 6). Thus, in continued optimization of the antiviral system, the protocol was adjusted to contain both 1:10 diluted stock MS2 bacteriophage suspension and solutions were not allowed to dry to evaporation.

Antiviral Capabilities of ClO₂ Gas on Cloth

Prior sets of studies determined that maximum antiviral potential was achieved when the stock solution was diluted in water (thus, reducing organic content) and when the material was inoculated and not allowed to dry before exposure to ClO₂ gas. Thus, the next phase of investigation examined the antiviral potential of ClO₂ gas on cloth (a porous substrate) under the same optimized treatment protocol. In testing three variables, concentration, exposure time length, and relative humidity, it was observed that all three influence the ability of ClO₂ to act as an antiviral agent (Table 2).

First, the effects of ClO₂ on cloth for 45 minutes at 200 PPM yielded an average of 95% viral reduction ($SD = 0.894$) and a 99.8 % reduction at 960 minutes (Figure 9). It was then determined that decreasing the concentration of ClO₂ gas to 40 PPM for 960 minutes altered the antiviral ability of ClO₂ gas to 24% ($SD = 0.021$) (Figure 10). Lastly, it was determined that increasing the relative humidity to that of $\geq 90\%$ and decreasing the concentration to 20 PPM increased the antiviral capabilities to a complete elimination of recoverable viral particles (Figure 11).

Table 1: Investigation of the effects of 200 ppm ClO₂ gas on MS2 inoculated stainless-steel coupons at room temperature and ≤ 64% relative humidity

Contaminated Substrate	Exposure Time (min)	Precent Reduction†
Dry Steel Coupon*	45	61%
Wet Steel Coupon**	45	87%
Wet Steel Coupon**	960	99.30%
Diluted Wet Steel Coupon***	45	99.80%
Diluted Wet Steel Coupon***	960	99.99%

* Contaminate left on metal coupon for 1 hour ± 15 minutes to allow contaminated surface to dry to evaporation.
 ** After inoculation, surface was immediately exposed to ClO₂ gas.
 *** MS2 bacteriophage solution was diluted with a 1:10 sterile water solution.
 † Average triplicate experiments

Table 2: Investigation of the effects of ClO₂ gas on MS2 impregnated cloth at room temperature and ≤ 64%- ≥90 % relative humidity

Exposure Time (min)	Concentration (PPM)	Relative Humidity (%)	Precent Reduction†	Log Reduction
45	200	≤ 64%	95%	1.9
960	200	≤ 64%	99.8%	2.9
960	40	≤ 64%	24%	0.1
960	20	≥ 90%	NRV* (≈100%)	≥ 7

* NRV = No Recoverable Viruses
 † Average duplicate experiments




Figure 7: Examination of the effects of wet versus dry viral suspension contaminated stainless steel surfaces exposed to 200 ppm ClO₂ at room temperature and ≤ 64% humidity for 45 minutes




Figure 8: Examination of the effects of 1:10 diluted MS2 bacteriophage stock solution with sterile H₂O prior to 200 ppm ClO₂ gas exposure for 45 minutes at room temperature and <64% humidity




Figure 9: Examination of the antiviral capabilities of ClO₂ on cloth impregnated with 1:10 diluted MS2 bacteriophage solution at 45 minutes and 960 minutes at room temperature and ≤ 64% relative humidity




Figure 10: Examination of the antiviral capabilities of ClO₂ on cloth impregnated with 1:10 diluted MS2 bacteriophage solution at varying concentrations of 200 PPM and 40 ppm ClO₂ Gas for 45 minutes at room temperature and ≤ 64% relative humidity




Figure 11: Examination of the antiviral capabilities of ClO₂ on cloth impregnated with 90% diluted MS2 bacteriophage solution at varying relative humidity conditions of ≤ 64% and ≥ 90% relative humidity (RH)

CHAPTER IV: DISCUSSION

Summary of Findings

In summary, it was found that several factors influence the ability of ClO₂ gas to act as an antiviral agent. Such factors that contribute to the gas's antiviral activity include the amount of organic content viral particles are suspended in, humidity, concentration of ClO₂ gas, and substrate. The optimal results of the antiviral capabilities of ClO₂ gas were found at $\geq 90\%$ relative humidity at room temperature with viral particles diluted in 90% sterile H₂O at 20 PPM of ClO₂ gas impregnated in cloth.

Initial findings revealed that percent reduction of viral particles is heavily dependent upon the organic concentration of the viral suspension. Since ClO₂ gas is a strong oxidizing compound, especially toward organic matter, it would be expected that the ClO₂ gas may be reduced to chlorine monoxide or to the chloride ion in the presence of higher levels of organic content (Hupperich et al. 2020; Wenk et al. 2013). The ability of ClO₂ to oxidize organic matter, implies that the viability of ClO₂ gas to act as an antiviral agent is reduced in the presence of organic content, thus, as the results show, reduction of the organic content has a direct effect on the ability of ClO₂ gas to inactivate the MS2 bacteriophage.

Additional findings revealed that ClO₂ gas had better ability acting as an antiviral agent on cloth rather than stainless steel metal coupons. Conceivably, this may be due to oxidation of the stainless steel by ClO₂ gas, although additional research is needed to come to such conclusion. One other potential theory is that ClO₂ gas penetrates deeper

into the cloth thus being exposed to more viral particles leading to an increase in viral deactivation.

Lastly, it was found that ClO₂ gas's ability to act as an antiviral agent is dependent upon the relative humidity of the environment it is deployed in. This finding stands in solidarity with other studies investigating ClO₂ gas's ability to act as an oxidizing agent towards biological agents at varying relative humidity (Park et al. 2018; Shirasaki et al. 2016). Studies suggest that this property of ClO₂ gas is due to the varying solubilities of ClO₂ gas at differing levels of humidity.

Implications

From the results, data supports the use of ClO₂ gas as an antiviral agent. The results yielded a sufficient disinfection of both cloth and metal to be considered an antiviral agent. This study offers insight into the various factors that must be considered when using ClO₂ gas as an antiviral agent and future obstacles that must be considered in pursuit of further ClO₂ gas antiviral research.

Such application of the findings include use of ClO₂ gas on metal and cloth as a potential antiviral agent, but also suggest that there may be a wider horizon of antiviral application for ClO₂ gas. Future studies should be aimed at investigation of ClO₂ gas on surfaces such as wood and plastics. It also may be of interest to expand research on the use of ClO₂ gas on bacteria' spores and mycobacteria which are known to be especially resistant to chemical inactivation.

Bibliography

- Aieta EM, Berg JD. 1986. A review of chlorine dioxide in drinking water treatment. *J Am Water Works Assoc.* 78(6):62–72. doi:10.1002/j.1551-8833.1986.tb05766.x.
- Alvarez ME, O'Brien RT. 1982. Mechanisms of inactivation of poliovirus by chlorine dioxide and iodine. *Applied and Environmental Microbiology.* 44(5):1064–1071. doi:10.1128/aem.44.5.1064-1071.1982.
- Bhawana J, Stubblefield JM, Newsome AL, Cahoon AB. 2014. Surface decontamination of plant tissue explants with chlorine dioxide gas. *In Vitro Cell Dev Biol - Plant.* 51(2):214–219. doi:10.1007/s11627-014-9659-4.
- Blancou J. 1995. Les méthodes de désinfection de l'antiquité à la fin du XVIIIe siècle. *Rev - Off Int Epizoot Copy.* 14(1):21–39. doi:10.20506/rst.14.1.831.
- Bruslind L. 2018 Feb 6. Introduction to Viruses. *Biology LibreTexts.*
[https://bio.libretexts.org/Bookshelves/Microbiology/Book%3A_Microbiology_\(Bruslind\)/08%3A_Introduction_to_Viruses.](https://bio.libretexts.org/Bookshelves/Microbiology/Book%3A_Microbiology_(Bruslind)/08%3A_Introduction_to_Viruses)
- Clokier MRJ, Millard AD, Letarov AV, Heaphy S. 2011. Phages in nature. *Bacteriophage.* 1(1):31–45. doi:10.4161/bact.1.1.14942.
[https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3109452/.](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3109452/)
- Cowan M, Smith H, Lusk J. 2019. *Microbiology fundamentals: a clinical approach.* 3rd ed. New York, NY: McGraw-Hill Education.

- Gaustad JW, McDuff CR, Hatcher HJ. 1974. Test method for the evaluation of virucidal efficacy of three common liquid surface disinfectants on a simulated environmental surface. *Appl Microbiol.* 28(5):748–752. doi:10.1128/am.28.5.748-752.1974.
- Gormsen M. 2016. The effectiveness of chlorine dioxide in inactivating influenza virus [Undergraduate Thesis]. [University Honors College, Middle Tennessee State University]. <https://jewlscholar.mtsu.edu/items/559317eb-7ea0-4d6c-9a61-42fd115d7c>.
- Hugo WB. 1991. A brief history of heat and chemical preservation and disinfection. *J Appl Microbiol.* 71(1):9–18. doi:10.1111/j.1365-2672.1991.tb04657.x.
- Hupperich K, Mutke XAM, Abdighahroudi MS, Jütte M, Schmidt TC, Lutze HV. 2020. Reaction of chlorine dioxide with organic matter – formation of inorganic products. *Environ Sci Water Res Technol.* 6(9):2597–2606. doi:10.1039/d0ew00408a.
- Isomoto H, Urata M, Kawazoe K, Matsuda J, Nishi Y, Wada A, Ohnita K, Hirakata Y, Matsuo N, Inoue K, et al. 2006. Endoscope disinfection using chlorine dioxide in an automated washer-disinfector. *J Hosp Infect.* 63(3):298–305. doi:10.1016/j.jhin.2006.01.020.
- Johns Hopkins University. 2020. COVID-19 dashboard. Johns Hopkins Coronavirus Resource Center; [updated 2021 Nov 14; accessed 2021 Nov 14]. <https://coronavirus.jhu.edu/map.html>.

- Kály-Kullai K, Wittmann M, Noszticzius Z, Rosivall L. 2020. Can chlorine dioxide prevent the spreading of coronavirus or other viral infections? Medical hypotheses. *Physiol Int.* 107(1):1–11. doi:10.1556/2060.2020.00015.
- Keen EC. 2015. A century of phage research: Bacteriophages and the shaping of modern biology. *BioEssays.* 37(1):6–9. doi:10.1002/bies.201400152.
- Lenntech. 2021. Disinfectants chlorine dioxide. Water Treatment Solutions. [accessed 2021 Nov 16]. <https://www.lenntech.de/chlorine-dioxide.htm>.
- Lin Q, Lim JYC, Xue K, Yew PYM, Owh C, Chee PL, Loh XJ. 2020. Sanitizing agents for virus inactivation and disinfection. *View.* (e16). <https://doi.org/10.1002/viw2.16>.
- Medawar P, Medawar J. 1985. Aristotle to zoos: a philosophical dictionary of biology. Cambridge (M.A.): Harvard University Press. 320 p.
- Newsome A, Stubblefield J. 2012. Novel disinfection applications using a portable chlorine dioxide gas generation system. In: EPA Decontamination Research and Development Conference. Vol. 600/R/12/557. Raleigh, NC. p. 61–62.
- Newsome A, DuBois JD, Tenney JD. 2009. Disinfection of football protective equipment using chlorine dioxide produced by the ICA TriNova system. *BMC Public Health.* 9(1). doi:10.1186/1471-2458-9-326.

- Noszticzius Z, Wittmann M, Kály-Kullai K, Beregvári Z, Kiss I, Rosivall L, Szegedi J. 2013. Chlorine dioxide is a size-selective antimicrobial agent. Schlievert P, editor. PLoS ONE. 8(11):1–10. doi:10.1371/journal.pone.0079157.
- Ogata N. 2012. Inactivation of influenza virus haemagglutinin by chlorine dioxide: oxidation of the conserved tryptophan 153 residue in the receptor-binding site. J Gen Virol. 93(12):2558–2563. doi:10.1099/vir.0.044263-0.
- Ogata N, Sakasegawa M, Miura T, Shibata T, Takigawa Y, Taura K, Taguchi K, Matsubara K, Nakahara K, Kato D, et al. 2016. Inactivation of airborne bacteria and viruses using extremely low concentrations of chlorine dioxide gas. Pharmacology. 97(5-6):301–306. doi:10.1159/000444503.
- Ogata N, Shibata T. 2008. Protective effect of low-concentration chlorine dioxide gas against influenza A virus infection. J Gen Virol. 89(1):60–67. doi:10.1099/vir.0.83393-0.
- Oldstone M. 2010. Viruses, plagues, and history: past, present, and future. New York (N.Y.): Oxford University Press. 383 p.
- Oudejans L, Ratliff K. 2021. COVID-19: UVC Devices and Methods for Surface Disinfection. EPA Office of Research and Development Homeland Security Research. https://cfpub.epa.gov/si/si_public_record_report.cfm?dirEntryId=350664&Lab=CESER&personid=23379&role=Any.

- Park SH, Kim WJ, Kang DH. 2018. Effect of relative humidity on inactivation of foodborne pathogens using chlorine dioxide gas and its residues on tomatoes. *Appl Microbiol.* 67(2):154–160. doi:10.1111/lam.13002.
- Pellett PE, Mitra S, Holland TC. 2014. Basics of virology. *Handb Clin Neurol.* 123(2014):45–66. doi:10.1016/b978-0-444-53488-0.00002-x.
- Pressbooks. 2014. The viral life cycle microbiology. Lumen Learning. [accessed 2021 Nov 23]. <https://courses.lumenlearning.com/microbiology/chapter/the-viral-life-cycle/>.
- Rabenau HF, Schwebke I, Blümel J, Eggers M, Glebe D, Rapp I, Sauerbrei A, Steinmann E, Steinmann J, Willkommen H, et al. 2020. Guideline for testing chemical disinfectants regarding their virucidal activity within the field of human medicine. *Bundesgesundheitsblatt - Gesundheitsforsch - Gesundheitsschutz.* 63(5):645–655. doi:10.1007/s00103-020-03115-w.
- Rastogi VK, Ryan SP, Wallace L, Smith LS, Shah SS, Martin GB. 2010. Systematic evaluation of the efficacy of chlorine dioxide in decontamination of building interior surfaces contaminated with anthrax spores. *Appl Environ Microbiol.* 76(10):3343–3351. doi:10.1128/AEM.02668-09.
- Ratliff C, Oudejans L. 2021. U.S. EPA research webinar - COVID-19: UV-C devices & methods for surface disinfection [PowerPoint slides]. Research Triangle Park (N.C.): Environmental Protection Agency.

https://cfpub.epa.gov/si/si_public_record_report.cfm?dirEntryId=350664&Lab=CESER&personid=23379&role=Any.

Sanekata T, Fukuda T, Miura T, Morino H, Lee C, Maeda K, Araki K, Otake T, Kawahata T, Shibata T. 2010. Evaluation of the antiviral activity of chlorine dioxide and sodium hypochlorite against feline calicivirus, human influenza virus, measles virus, canine distemper virus, human herpesvirus, human adenovirus, canine adenovirus and canine parvovirus. *Biocontrol Sci.* 15(2):45–49. doi:10.4265/bio.15.45.

Schwartz E. 2019. How to Calculate Virus Titers. *Sciencing*. [accessed 2021 Nov 27]. <https://sciencing.com/how-6610893-calculate-virus-titers.html>.

Shirasaki Y, Matsuura A, Uekusa M, Ito Y, Hayashi T. 2016. A study of the properties of chlorine dioxide gas as a fumigant. *Exp Anim.* 65(3):303–310. doi:10.1538/expanim.15-0092.

Smith DJ, Ernst W, Giddings JM. 2014. Distribution and chemical fate of ³⁶Cl-chlorine dioxide gas during the fumigation of tomatoes and cantaloupe. *J Agric Food Chem.* 62(48):11756–11766. doi:10.1021/jf504097p.

Stubblefield JM, Newsome AL. 2015. Potential biodefense model applications for portable chlorine dioxide gas production. *Health Secur.* 13(1):20–28. doi:10.1089/hs.2014.0017.

- Virox Technologies. History of disinfection. History of Disinfection - Virox Technologies Inc. [accessed 2021 Nov 17]. <https://virox.com/history-of-disinfection/>.
- Wenk J, Aeschbacher M, Salhi E, Canonica S, von Gunten U, Sander M. 2013. Chemical oxidation of dissolved organic matter by chlorine dioxide, chlorine, and ozone: effects on its optical and antioxidant properties. *Environ Sci Technol.* 47(19):11147–11156. doi:10.1021/es402516b.
- Woolwine JD, Gerberding JL. 1995. Effect of testing method on apparent activities of antiviral disinfectants and antiseptics. *Antimicrob Agents Chemother.* 39(4):921–923. doi:10.1128/aac.39.4.921.
- Wyrzykowska-Ceradini B, Calfee MW, Touati A, Wood J, Mickelsen RL, Miller L, Colby M, Slone C, Gatchalian NG, Pongur SG, et al. 2019. The use of bacteriophage MS2 for the development and application of a virucide decontamination test method for porous and heavily soiled surfaces. *J Appl Microbiol.* 127(5):1315–1326. doi:10.1111/jam.14406.